

Spherical Gravitational Collapse of Annihilating Dark Matter and the Minimum Mass of CDM Black Holes

R. Ali Vanderveld and Ira Wasserman

Center for Radiophysics and Space Research, Cornell University, Ithaca NY 14853

Spherical gravitational collapse of a cold gas of annihilating particles involves a competition between two density dependent rates: the free-fall rate $\propto \sqrt{\rho}$ and the (s-wave) annihilation rate $\propto \rho$. Thus, there is a critical density ρ_{ann} above which annihilation proceeds faster than free fall. Gravitational collapse of a cloud of (initial) mass M to a black hole is only possible if $\rho_{\text{BH}} \equiv 3M/4\pi(2GM)^3 = 3/32\pi G^3 M^2 \lesssim \rho_{\text{ann}}$, which implies that $M \gtrsim M_{\text{ann}} \equiv (3/32\pi G^3 \rho_{\text{ann}})^{1/2}$. We compute the collapse and annihilation of cold, spherical gas clouds with either a homogeneous density profile or a specific radially-dependent profile that would lead to self-similar collapse if there were no annihilation. These calculations verify that there exists a lower bound to the range of possible black hole masses that can form in cold dark matter (CDM) collapse. For a particle mass m and freeze-out temperature $T_f = m/x_f$, the minimum black hole mass is $M_{\text{ann}} \approx 10^{10} M_{\odot} \times (x_f \sqrt{g_{\star}}/100\Omega_{\text{CDM}} h^2 g_{\star} s m(\text{Gev}))$, where $g_{\star}$ and $g_{\star}$ are degeneracy factors. Moreover, the formation of a black hole is accompanied by the annihilation of about M_{ann} of the original mass. Most of the annihilated mass is released in a burst lasting a time $\sim GM_{\text{BH}}$, where M_{BH} is the black hole mass when collapse first occurs. A total annihilation luminosity $\sim 10^{59} \text{ erg s}^{-1}$ could easily be reached, lasting a time ~ 100 seconds. The absence of astronomical observations of such spectacular events suggests either: (i) the branching ratio for CDM annihilation to e^+e^- pairs or quarks $\lesssim 10^{-10}$, while the branching ratio to $\nu\bar{\nu}$ is $\lesssim 10^{-5}$; or (ii) CDM is not made of annihilating particles, but may be in some non-annihilating form, such as axions; or (iii) CDM black holes never form.

PACS numbers: 95.35.+d, 97.60.Lf

I. INTRODUCTION

Non-baryonic matter dominates the matter content of the Universe today, with a density parameter $\Omega_{\text{CDM}} h^2 \approx 0.15$; the rest of the energy density is primarily in the form of dark energy [1, 2]. In the favored cold dark matter (CDM) scenario based on weakly interacting massive particles (WIMPS), the dark matter is in the form of a gas of particles that is very cold today, with an abundance that was lowered substantially by annihilation when the temperature of the Universe was $\gtrsim m/x_f$, for a CDM particle mass m freezing out at temperature $T_f = m/x_f \sim 0.1m$ [3, 4, 5]. The Λ CDM picture provides acceptable fits to the temperature and E-polarization fluctuations of the cosmic microwave background (CMB) radiation [6, 7], as well as to the large scale structure of the Universe [8]. On smaller scales, CDM halos seem to form cuspy cores [9, 10]; dark matter annihilation has been proposed as one means of smoothing out the cusps [11, 12]. CDM clumps are found to form in large scale structure simulations and to agglomerate to form halos [13]; annihilation within these clumps would lead to an enhanced flux of cosmic ray products at Earth [14, 15].

In this paper, we investigate a related question: How does annihilation alter spherically symmetric collapse of CDM? We consider, for the most part, the collapse of CDM clouds with zero or very small internal energy. In the absence of annihilation, the inevitable result would be complete gravitational collapse to a black hole.

However, to form a black hole, a collapsing sphere of mass M must reach a mass density of order $\rho_{\text{BH}} = 3M/4\pi(2GM)^3 = 3/32\pi G^3 M^2$. When (s-wave) annihilation is included, the rate of CDM annihilation at a density ρ is

$$\Gamma_{\text{ann}} = \frac{\rho \langle \sigma v \rangle_{\text{ann}}}{m}, \quad (1)$$

where $\langle \sigma v \rangle_{\text{ann}}$ is the annihilation rate coefficient and m is the CDM particle mass. For comparison, the free fall rate at high densities is approximately

$$H = \sqrt{\frac{8\pi G \rho}{3}}; \quad (2)$$

then $\Gamma_{\text{ann}} = H$ at a density

$$\rho_{\text{ann}} = \frac{8\pi G m^2}{3 (\langle \sigma v \rangle_{\text{ann}})^2}. \quad (3)$$

For $\rho \lesssim \rho_{\text{ann}}$, free fall proceeds faster than annihilation, and vice-versa. Therefore, black hole formation is only possible for a mass M provided that $\rho_{\text{BH}} \lesssim \rho_{\text{ann}}$ or

$$M \gtrsim \frac{3\langle\sigma v\rangle_{\text{ann}}}{16\pi G^2 m} \equiv M_{\text{ann}}. \quad (4)$$

The annihilation of CDM particles in the early Universe leads to a residual density parameter $\Omega_{\text{CDM}}h^2$ provided that

$$\langle\sigma v\rangle_{\text{ann}} \approx 8.6 \times 10^{-9} \text{ GeV}^{-2} \left(\frac{x_f}{100\Omega_{\text{CDM}}h^2 g_{*S}/\sqrt{g_*}} \right), \quad (5)$$

where g_{*S} and g_* are degeneracy factors for the entropy and mass density when the temperature of the Universe is $\sim T_f$, and

$$x_f \approx \ln C - \frac{1}{2} \ln(\ln C), \quad (6)$$

where $C = 0.038(g/\sqrt{g_*})m\langle\sigma v\rangle_{\text{ann}}/\sqrt{G}$, with g being the degeneracy factor for the CDM particles alone [16]. Substituting for $\langle\sigma v\rangle_{\text{ann}}$ in equation (4), we find

$$M_{\text{ann}} \approx \frac{10^{10} M_{\odot}}{m(\text{GeV})} \left(\frac{x_f}{100\Omega_{\text{CDM}}h^2 g_{*S}/\sqrt{g_*}} \right) \approx \frac{10^{10} M_{\odot}}{m(\text{GeV})} \left(\frac{x_f}{15g_{*S}/\sqrt{g_*}} \right), \quad (7)$$

where $m = m(\text{GeV}) \text{ GeV}$; furthermore, in our last result, we assume that $\Omega_{\text{CDM}}h^2 \approx 0.15$, as has been determined from analyses of the CMB and of large scale structure. Our estimate of M_{ann} depends on the CDM particle mass in several ways, since $M_{\text{ann}} \propto x_f \sqrt{g_*}/mg_{*S}$. In addition to the explicit inverse proportionality with m , there is a logarithmic dependence from x_f and a piecewise flat dependence via g_{*S} and g_* : for example, if $m = 50 \text{ GeV}$, we would have enough particle species for $g_* \approx g_{*S} \approx 50$, and $M_{\text{ann}} \approx 4 \times 10^7 M_{\odot}$ (taking $x_f \approx 20$ for this case). Plausibly, the value of the minimum black hole mass for CDM is comparable to the mass range deduced for black holes in the centers of galaxies [17]. It is remarkable (although perhaps fortuitous) that the value of M_{ann} is even remotely close to an astronomically significant value.

Spherical collapse is, of course, highly idealized. More realistically, we would expect a collapsing cloud to have substantial substructure as well as large scale shear flows induced by the tidal field of nearby fluctuations. However, we would still expect that CDM clouds that are, however improbably, nearly spherical and smooth would have the *best chance* to collapse to black holes. Thus, our lower bound on the CDM black hole mass ought to be robust. Moreover, shear enhances the collapse of a cloud (as is evident from the Raychauduri equation, e.g. [18]) and anisotropic collapse of a cloud with internal fluctuations could trigger substantial fragmentation (e.g. [19, 20] and refs. therein). It is possible that the calculations we present apply to late time collapse within a larger cloud that has fragmented considerably.

Section II presents calculations of the collapse of homogenous CDM spheres. Section III computes the collapse of a specific model with radial density dependence. In each case, we quantify when black holes may form, and when the collapsing mass will instead annihilate away. The detailed calculations are Newtonian, and assume instantaneous escape of the (relativistic) products of CDM annihilation. We expect that our main results for the critical mass for black hole formation, and for the mass-energy and luminosity released by annihilation, to remain valid within factors ~ 1 in a general relativity calculation that also accounts for the opacity of the infalling CDM to the annihilation products. In Section IV, we discuss our results, including a brief account of what might happen in the collapse of CDM with predominantly p-wave annihilation.

II. THE HOMOGENEOUS CASE

A. Equations of motion

Let us consider a uniform sphere of matter with time-dependent radius and mass, called $R(t)$ and $M(t)$, respectively; hence, the density at a time t is $\rho(t) = 3M(t)/4\pi R^3(t)$. We assume that the homogeneous sphere collapses under its own self gravity, but we also include pressure forces. To maintain homogeneity, we assume that

$$P(r, t) = P_0(t) \left[1 - \frac{r^2}{R^2(t)} \right] \quad (8)$$

at time t and Eulerian radius r ; we also assume that $P_0(t) = K\rho^\gamma(t)$, where K is independent of time. The Euler equation then becomes

$$\ddot{R} = \frac{2K(3M/4\pi)^{\gamma-1}}{R^{3\gamma-2}} - \frac{GM}{R^2}. \quad (9)$$

If $\gamma > 4/3$, then there exists a radius R_{ZA} at which $\ddot{R} = 0$. For a nonrelativistic gas we expect to have $\gamma = 5/3$, so in this case we have the zero acceleration radius

$$R_{ZA} = \frac{2K}{GM^{1/3}} \left(\frac{3}{4\pi} \right)^{2/3}. \quad (10)$$

This is a minimum radius at which the pressure force becomes as great as the gravitational force such that the collapse will “bounce” back outward. In the absence of annihilation, we also have the conserved energy

$$E = \frac{1}{2}\dot{R}^2 - \frac{GM}{R} + \left(\frac{3M}{4\pi} \right)^{2/3} \frac{K}{R^2}. \quad (11)$$

From this we see that the cloud will bounce at a radius $R = \frac{1}{2}R_{ZA}$ in the absence of annihilation.

We note that there may be three possible physical sources of “pressure”: (i) internal motions associated with small scale density inhomogeneities within a cloud; (ii) residual but small entropy of the CDM particles; (iii) trapped annihilation products.

Of these three sources, the first is potentially the most important, at least on large scales, but, as discussed in the introduction, we are focusing on fragments that have the best chance for collapse. Thus, we assume that our collapsing cloud is smooth; the larger scale motions associated with inhomogeneities of the parent system are assumed to be in the form of orbital motion of the subclumps into which it has fragmented by the time our smooth cloud collapses to very high density.

The residual entropy of the CDM is small and depends on the scattering cross section of thermalized particles in the early Universe with the CDM. Assuming a cross-section $f_t \langle \sigma v \rangle_{\text{ann}} (T/m)^2$, the rate of energy transfer from thermal particles to CDM is $\dot{U}_t = \zeta(3)g_t f_t \langle \sigma v \rangle_{\text{ann}} T^7 / \pi^2 m^3$ per CDM particle, where g_t is the spin weight of relativistic thermal particles that scatter from the CDM and f_t is a numerical factor of order one. The CDM therefore remains in “kinetic” thermal contact until $\Gamma_t = 2\dot{U}_t/3T = H = (8\pi^3 g_\star G T^4/90)^{1/2}$, which implies kinetic freeze-out at [4]

$$T_K \approx \frac{1.9g_\star^{1/8} G^{1/8} m^{3/4}}{g_t^{1/4} f_t^{1/4} (\langle \sigma v \rangle_{\text{ann}})^{1/4}} \approx 3.4 \times 10^{-3} \text{ GeV} \frac{g_\star^{1/4} [m(\text{GeV})]^{3/4}}{g_t^{1/4} f_t^{1/4}} \left(\frac{100\Omega_{\text{CDM}} h^2}{x_f} \right)^{1/4}; \quad (12)$$

the value of K associated with this residual thermal energy is then

$$\begin{aligned} K_{\text{CDM}} &= \frac{T_K}{m[\rho(T_K)]^{2/3}} \\ &\approx 4.9 \times 10^8 \text{ GeV}^{-8/3} \frac{g_t^{1/4} f_t^{1/4}}{[m(\text{GeV})]^{7/4} g_\star^{1/4} g_\star^{2/3}} \left(\frac{x_f}{100\Omega_{\text{CDM}} h^2} \right)^{1/4} \left(\frac{0.15}{\Omega_{\text{CDM}} h^2} \right)^{2/3}, \end{aligned} \quad (13)$$

where $g_\star$ is the degeneracy factor for the entropy at temperature T_K . To put this in context, the CDM phase space density is $\sim K_{\text{CDM}}^{-3/2} m^{-4} \sim 10^{-13} [m(\text{GeV})]^{-11/8}$. The CDM phase space density and, therefore, K_{CDM} are conserved as long as the CDM particles are collisionless.

The third source of pressure would be from trapped products of the annihilating dark matter. For the most part, we shall assume that these annihilation products escape without being trapped, and this further assumption will be discussed below. (We shall also re-examine this assumption semi-quantitatively in our brief discussion of the collapse of CDM that annihilates predominantly via p-wave interactions.) Any products that are trapped should be extremely relativistic and therefore would form a $\gamma = 4/3$ gas. At fixed K , a $\gamma = 4/3$ gas either stops the collapse at all times, or not at all.

Annihilation implies a mass loss rate

$$\dot{M} = - \frac{3\langle \sigma v \rangle_{\text{ann}} M^2}{4\pi m R^3}. \quad (14)$$

Here, we assume that the annihilation products escape the collapsing cloud completely. This and equation (9) give us our system of coupled differential equations which we must solve. These look much simpler once nondimensionalized

with a change to new mass, radius and time units. Let the mass unit be the initial mass M_0 such that the new mass variable is $\hat{M}(t) = M(t)/M_0$. Then introduce a radius unit R_u and a time unit $\sqrt{R_u^3/GM_0}$ for the new variables $\hat{R}$ and τ , respectively. Then the above equations become

$$\frac{d^2 \hat{R}}{d\tau^2} = \frac{R_{ZA} \hat{M}^{2/3}}{R_u \hat{R}^3} - \frac{\hat{M}}{\hat{R}^2} \quad (15)$$

$$\frac{d\hat{M}}{d\tau} = -\frac{3\langle\sigma v\rangle_{\text{ann}}}{4\pi m G} \sqrt{\frac{GM_0}{R_u^3}} \frac{\hat{M}^2}{\hat{R}^3} \equiv -\left(\frac{R_{\text{ann}}}{R_u}\right)^{3/2} \frac{\hat{M}^2}{\hat{R}^3}; \quad (16)$$

here the characteristic radius at which annihilation becomes important is defined to be

$$\begin{aligned} R_{\text{ann}} &\equiv \frac{3\langle\sigma v\rangle_{\text{ann}}}{4^{4/3}\pi G m} \left(\frac{M_0}{M_{\text{ann}}}\right)^{1/3} \\ &= \frac{3.8 \times 10^{15} \text{ cm}}{m(\text{GeV})} \left(\frac{x_f}{100\Omega_{\text{CDM}} h^2 g_{\star S}/\sqrt{g_{\star}}}\right) \left(\frac{M_0}{M_{\text{ann}}}\right)^{1/3}. \end{aligned} \quad (17)$$

We choose $R_u = \sqrt{R_{\text{ann}}^2 + R_{\text{ZA}}^2}$; letting $\kappa = R_{\text{ZA}}/R_u$ and $R_{\text{ann}} = R_u \sqrt{1 - \kappa^2}$, the equations describing the collapse become

$$\begin{aligned} \frac{d^2 \hat{R}}{d\tau^2} &= \frac{\kappa \hat{M}^{2/3}}{\hat{R}^3} - \frac{\hat{M}}{\hat{R}^2} \\ \frac{d\hat{M}}{d\tau} &= -\frac{(1 - \kappa^2)^{3/4} \hat{M}^2}{\hat{R}^3}. \end{aligned} \quad (18)$$

The characteristic timescale for the cloud's evolution is

$$\begin{aligned} t_u &= \sqrt{\frac{R_u^3}{GM_0}} = \frac{3\langle\sigma v\rangle_{\text{ann}}}{4\pi G m} \left(\frac{R_u}{R_{\text{ann}}}\right)^{3/2} \\ &\approx \frac{2.0 \times 10^5 \text{ sec}}{m(\text{GeV})(1 - \kappa^2)^{3/4}} \left(\frac{x_f}{100\Omega_{\text{CDM}} h^2 g_{\star S}/\sqrt{g_{\star}}}\right). \end{aligned} \quad (19)$$

For cases where pressure forces are unimportant, $\kappa \rightarrow 0$; for $K = K_{\text{CDM}}$

$$\frac{R_{\text{ZA}}}{R_{\text{ann}}} = \frac{\kappa}{\sqrt{1 - \kappa^2}} \approx \frac{1.3 \times 10^{-5} g_t^{1/4} f_t^{1/4} g_{\star S}^{13/12}}{g_{\star SK}^{2/3} g_{\star}^{2/3} [m(\text{GeV})]^{5/12}} \left(\frac{100\Omega_{\text{CDM}} h^2}{x_f}\right)^{13/12} \left(\frac{0.15}{\Omega_{\text{CDM}} h^2}\right)^{2/3} \left(\frac{M_{\text{ann}}}{M_0}\right)^{2/3}, \quad (20)$$

which is indeed very small.

The solution of equations (18) depends on the value of κ and the initial conditions. Well before annihilation becomes important, $\hat{M}$ remains constant, and

$$\left(\frac{d\hat{R}}{d\tau}\right)^2 = -\frac{\kappa}{\hat{R}^2} + \frac{2}{\hat{R}} + E. \quad (21)$$

For $E = 0$, we can integrate to find

$$\sqrt{2}|\tau - \tau_{\text{bounce}}| = \frac{2}{3} \left(\hat{R} - \frac{\kappa}{2}\right)^{3/2} + \kappa \left(\hat{R} - \frac{\kappa}{2}\right)^{1/2}, \quad (22)$$

where the bounce happens when $\tau = \tau_{\text{bounce}}$. We use equation (22) to estimate the initial time for our numerical solutions.

As for the initial conditions, $\hat{M}(\tau) = 1$ at the start and $\hat{R}(\tau)$ should ideally fall from a very large distance, where $d\hat{R}/d\tau \approx 0$. However, since annihilation occurs only at small radii $\hat{R} \lesssim 1$, we can initialize numerical calculations to excellent accuracy by setting $\hat{M}_0 = 1$ as long as the initial value $\hat{R}_0$ is sufficiently large. We can then find the corresponding initial infall velocity by assuming previously negligible annihilation; in this limit, we can use the above energy conservation equation to find

$$\left(\frac{d\hat{R}}{d\tau}\right)_0 = -\sqrt{\frac{2}{\hat{R}_0} - \frac{\kappa}{\hat{R}_0^2}}, \quad (23)$$

which will be used in the numerical work described below.

B. Numerical calculation

Equations (18) were solved using a fourth-order Runge-Kutta routine [21]. The initial conditions for the mass, radius and velocity were set as discussed above; $\hat{R}_0 = 10$ turned out to be sufficiently large. The results, for different κ values, are displayed in Figure 1. Note how the minimum $\hat{R}$ and the final $\hat{M}$ values both appear to scale with κ . Also, we see that the $\kappa = 0$ case yields a power-law relationship between $\hat{M}$ and $\hat{R}$ for small $\hat{M}$ and $\hat{R}$. We can understand these relationships analytically by looking at the small κ and small $\hat{M}$ limit of the equations of motion.




FIG. 1: The top two plots show $\log(\hat{R})$ and $\log(\hat{M})$ versus $\log(\tau)$, for a time interval near the bounce and for $\kappa = 0.03, 0.01, 0.003$ and 0. The bottom plot is $\log(\hat{M})$ versus $\log(\hat{R})$ for the same four values of κ . The initial radius was $\hat{R}_0 = 10$.

Let us change the independent variable in our equations from τ to $x = 1/\hat{M}$ while defining the new dependent variable $y = \hat{R}^{-2}$. Rearranging the equations of motion yields

$$\sqrt{y} \frac{d^2 y}{dx^2} = \frac{2}{x}. \quad (24)$$

This form is more useful for finding the behavior of the solution when $\hat{M} \rightarrow 0$, which is equivalent to $x \rightarrow \infty$. In this case, let us look for a solution of the form

$$y = Ax + Bx^a + \dots \quad (25)$$

where the solution will only be sensible if the second term is much smaller than the first. By inserting this into equation (25) we find that

$$y \simeq Ax - 8\sqrt{\frac{x}{A}} + \dots \quad (26)$$

for large x ; the coefficient A is not determined by this argument. Note that, by returning to the original variables, this implies that $\hat{M}/\hat{R}^2$ approaches a constant as the sphere collapses. Putting κ back into the equations, but still requiring it to be small, we can take the $\hat{R} \rightarrow 0$ limit in order to predict a final mass $\hat{M}_f \propto \kappa^3$ and a minimum radius $\frac{1}{2}\hat{R}_{ZA} \propto \kappa^{3/2}$. These scalings agree with the numerical solutions.

The asymptotic scaling $\hat{M}/\hat{R}^2 \rightarrow \text{constant}$ for an annihilating cloud implies that $dR/dt \rightarrow \text{constant}$ at late times. Before annihilation becomes dominant, the cloud collapses in free fall; this phase ends when the cloud attains a density $\sim \rho_{\text{ann}}$. At that time, the cloud collapses at a rate $dR/dt \approx v_{\text{ann}} \equiv (M_0/M_{\text{ann}})^{1/3}$; subsequently, the cloud loses

mass, its self-acceleration diminishes, and it continues to collapse at roughly this speed. In doing so, its density is determined by the condition that the rates of annihilation and collapse remain about the same, i.e.

$$\frac{\rho\langle\sigma v\rangle_{\text{ann}}}{m} = \frac{3M\langle\sigma v\rangle_{\text{ann}}}{4\pi R^3 m} \sim \frac{v_{\text{ann}}}{R}, \quad (27)$$

which implies

$$\frac{M}{R^2} \sim \frac{4\pi m v_{\text{ann}}}{3\langle\sigma v\rangle_{\text{ann}}}. \quad (28)$$

Equation (28) corresponds to $\hat{M}/\hat{R}^2 = A \sim 4^{1/3}$.

At this point, we pause to consider the circumstances under which our assumption that annihilation products escape promptly is justified. Let us suppose that, when first produced with individual particle energy m , the relativistic annihilation products interact with the CDM particles with a cross section $f_s\langle\sigma v\rangle_{\text{ann}}$, where we expect $f_s \sim 1$. Then the optical depth of the collapsing cloud to the relativistic annihilation products is $\tau = f_s\rho\langle\sigma v\rangle_{\text{ann}}R/m$. The density at which the cloud first becomes opaque to these products is

$$\rho_1 = \rho_{\text{ann}} f_s^{-3/2} \left(\frac{M_{\text{ann}}}{M} \right)^{1/2} = \rho_{\text{BH}} f_s^{-3/2} \left(\frac{M}{M_{\text{ann}}} \right)^{3/2}. \quad (29)$$

Thus, we see that the cloud remains transparent to the annihilation products for cloud masses well above and well below M_{ann} . For cloud masses $\approx M_{\text{ann}}$, the optical depth rises to $\tau \sim 1$ just as the cloud reaches $\rho_{\text{ann}} \sim \rho_{\text{BH}}$. For these cases, the collapse of the cloud could be altered slightly, delaying the final outcome – total collapse or total annihilation – for a short time, but probably not for longer than the characteristic free fall or annihilation time.

C. Black hole formation

In order to analyze the black hole formation process, define the dimensionless “horizon parameter”

$$h(\tau) = \frac{\hat{M}(\tau)}{\hat{R}(\tau)} = \frac{M(\tau)}{R(\tau)} \frac{R_u}{M_0}, \quad (30)$$

which increases during the infall phase, reaches a maximum at the bounce and then decreases during the outflow phase. For $\kappa \neq 0$, this simplifies to

$$h(\tau) = \frac{2K}{G\kappa} \left(\frac{3}{4\pi M_0^2} \right)^{2/3} \frac{M(\tau)}{R(\tau)}, \quad (31)$$

and for $\kappa = 0$ we get

$$h(\tau) = \left(\frac{3\langle\sigma v\rangle_{\text{ann}}}{4\pi m\sqrt{G}M_0} \right)^{2/3} \frac{M(\tau)}{R(\tau)}. \quad (32)$$

We can easily find the maximum value of $h(\tau)$ for any particular value of κ , which ranges from $h_{\text{max}}(\kappa = 0) \approx 0.8$ to $h_{\text{max}}(\kappa = 1) = 2$. This is shown in Figure 2. In order for a black hole to form, we would require that

$$\frac{2GM}{R} > 1 \quad (33)$$

and this corresponds to a restriction on M_0 . For $\kappa \neq 0$, we require

$$M_0 \geq \sqrt{\frac{3}{4\pi}} \left(\frac{K}{\kappa G^2 h_{\text{max}}(\kappa)} \right)^{3/4} \quad (34)$$

for black hole formation, and similarly for $\kappa = 0$ we would require

$$M_0 \geq \frac{3\langle\sigma v\rangle_{\text{ann}}}{4\pi m G^2} \left(\frac{1}{2h_{\text{max}}(0)} \right)^{3/2} = \frac{\sqrt{2}M_{\text{ann}}}{[h_{\text{max}}(0)]^{3/2}}. \quad (35)$$

Therefore there is a definite minimum initial mass $\sim (1 - 2)M_{\text{ann}}$ required in order for the self-gravity of the system to overcome the annihilation and pressure.




FIG. 2: The top plot shows $h_{\max}$ versus κ and the bottom plot shows the values of $\hat{M}$ and $\hat{R}$ at this value of h , also versus κ . Initial radius $\hat{R}_0 = 10$.

III. A RADIAL DENSITY PROFILE

A. Equations of motion

A purely collisionless gas cloud, defined to have zero pressure and to be non-annihilating, collapsing under its own self-gravity can be visualized as a set of mass shells in free fall. We can label each shell by the mass it encloses, m_s , such that its radius at time t is $r(m_s, t)$. As long as no two shells cross, m_s remains unchanged and can be used as a comoving coordinate; this remains true when annihilation is included if we choose m_s to be the mass enclosed long before annihilation becomes important. For purely collisionless infall, the motion of an individual mass shell is determined by its conserved energy $E(m_s)$ and its “bang time” $t_0(m_s)$, which is defined by $r(m_s, t_0(m_s)) = 0$.

Self-similar solutions for the collapse of a collisionless cloud have been constructed by Penston [22] and Lynden-Bell and Lemos [23], hereafter LL. These models correspond to $E(m_s) = 0$ and a power-law $t_0(m_s) = t_0 + Bm_s^b$; the exact solution is then

$$r(m_s, t) = \left(\frac{9Gm_s}{2} \right)^{1/3} (t_0(m_s) - t)^{2/3} = \left(\frac{9Gm_s}{2} \right)^{1/3} (Bm_s^b + t_0 - t)^{2/3}. \quad (36)$$

The density and velocity profiles corresponding to this solution are

$$\begin{aligned} \rho(m_s, t) &= \frac{1}{4\pi[r(m_s, t)]^2(\partial r(m_s, t)/\partial m)} = \frac{1}{6\pi G(Bm_s^b + t_0 - t)[Bm_s^b(1 + 2b) + t_0 - t]} \\ v(m_s, t) &= -\sqrt{\frac{2Gm_s}{r(m_s, t)}} = -\left(\frac{4Gm_s}{3(Bm_s^b + t_0 - t)} \right)^{1/3} \end{aligned} \quad (37)$$

where the density profile differs from LL equation (2.9), which has a typo. Equations (36) and (37) describe a collapse profile consisting of a *shrinking* core of mass

$$m_c \sim [(t_0 - t)/B]^{1/b} \quad (38)$$

and radius

$$r_c \sim (18G)^{1/3} B^{-1/3b} (t_0 - t)^{(1+2b)/3b}. \quad (39)$$

Within the core, the density is nearly uniform, $\rho \sim \rho_c = 1/6\pi G(t_0 - t)^2$, and the velocity profile is nearly linear, $v(r, t) \sim -r\sqrt{8\pi G\rho_c}$. This is surrounded by a halo, in which the density and velocity profiles are approximately

powerlaws:

$$\begin{aligned}\rho &\sim \rho_c(m_c/m_s)^{2b} \sim \rho_c(r_c/r)^{6b/(1+2b)} \\ v &\sim v_c(m_c/m)^{(b-1)/3} \sim v_c(r_c/r)^{(b-1)/(1+2b)} .\end{aligned}\quad (40)$$

In the LL solution, the core density rises to infinity as $t \rightarrow t_0$, so black hole formation is inevitable in this model.

Our goal is to numerically compute the solution to this problem in the presence of annihilation. As in the preceding section, we will choose initial conditions such that annihilation will have been negligible previously, making our use of the LL solution a physically meaningful choice. The above solution has its largest density at the center, so we must specify the central density to be small enough such that it is sufficiently below the critical density specified in equation (3). The central density at the initial time $t = 0$ is

$$\rho(m_s = 0, t = 0) = \frac{1}{6\pi G t_0^2} . \quad (41)$$

For our numerical simulations, we require that

$$\rho_{ann} = \frac{8\pi\alpha}{3}\rho(m_s = 0, t = 0) ; \quad (42)$$

this puts a constraint on the constant t_0 :

$$t_0 = \sqrt{\frac{\alpha}{6\pi} \frac{\langle \sigma v \rangle_{ann}}{Gm}} , \quad (43)$$

where we choose the constant factor α such that the initial central density is sufficiently small. We chose $\alpha = 3/4\pi \times 10^6$, which starts the collapse at such an early time that the central density is approximately a million times smaller than ρ_{ann} , and therefore the collapse is still in freefall with utterly negligible annihilation. In this way, we will be justified in initializing our simulation with data from the LL solution.

In order to determine the other constant B for our initial conditions, we can now specify an initial density contrast between the center and the outer edge of the gas cloud; specifying this contrast is also necessary for numerical purposes because the LL solution extends off to infinity. Let the central density be a factor of β_i larger than that of the edge at the initial time; this assumption, along with density equation (37), tells us that we must have

$$B = \frac{\gamma t_0}{M_0^b} = \sqrt{\frac{\alpha}{6\pi} \frac{\gamma \langle \sigma v \rangle_{ann}}{Gm M_0^b}} , \quad (44)$$

where we have once again chosen to call the total initial mass M_0 and γ is the constant

$$\gamma = \frac{\sqrt{(b+1)^2 + (2b+1)(\beta_i - 1)} - (b+1)}{2b+1} . \quad (45)$$

The above conditions give us the constants and initial data necessary for a numerical computation.

Although we are using the initial contrast β_i for our initial data, the physically important density contrast is what we will call β_{ann} , which is the contrast when the center begins annihilating. This is the parameter which will become important when discussing black hole formation for this system. We can write β_i as a function of β_{ann} and α . First, define the annihilation time t_{ann} to be the time at which the central density would reach the annihilation density in the LL model:

$$\rho_{ann} = \frac{1}{6\pi G(t_0 - t_{ann})^2} . \quad (46)$$

Then $\beta_{ann} = \rho_{ann}/\rho(M_0, t_{ann})$, which is

$$\beta_{ann} = \left(BM_0^b \sqrt{6\pi G \rho_{ann}} + 1 \right) \left[(1+2b) BM_0^b \sqrt{6\pi G \rho_{ann}} + 1 \right] \quad (47)$$

and we can solve this equation for $BM_0^b \sqrt{6\pi G \rho_{ann}}$ to get

$$BM_0^b \sqrt{6\pi G \rho_{ann}} = -\frac{1+b}{1+2b} + \sqrt{\frac{\beta_{ann} - 1}{1+2b} + \left(\frac{1+b}{1+2b} \right)^2} . \quad (48)$$

The initial density contrast is

$$\beta_i = \left(\frac{BM_0^b}{t_0} + 1 \right) \left(\frac{(1+2b)BM_0^b}{t_0} + 1 \right) \quad (49)$$

which can be rewritten as

$$\beta_i = 1 + 2(1+b) \left(\frac{BM_0^b \sqrt{6\pi G \rho_{ann}}}{t_0 \sqrt{6\pi G \rho_{ann}}} \right) + (1+2b) \left(\frac{BM_0^b \sqrt{6\pi G \rho_{ann}}}{t_0 \sqrt{6\pi G \rho_{ann}}} \right)^2 ; \quad (50)$$

plugging in what we know from the equations above, we find

$$\begin{aligned} \beta_i = & 1 + 2(1+b) \sqrt{\frac{3}{8\pi\alpha}} \left[\sqrt{\frac{\beta_{ann}-1}{1+2b} + \left(\frac{1+b}{1+2b} \right)^2} - \frac{1+b}{1+2b} \right] \\ & + (1+2b) \left(\frac{3}{8\pi\alpha} \right) \left[\sqrt{\frac{\beta_{ann}-1}{1+2b} + \left(\frac{1+b}{1+2b} \right)^2} - \frac{1+b}{1+2b} \right]^2 . \end{aligned} \quad (51)$$

We will use β_i and α in the calculation of initial conditions, but whether or not a black hole forms will be sensitive only to our choice of β_{ann} .

The addition of annihilation requires that we add the mass loss equation

$$\frac{\partial \mu(m_s, t)}{\partial t} = - \frac{\langle \sigma v \rangle_{ann}}{m} \rho(m_s, t) \mu(m_s, t) \quad (52)$$

where the comoving variable m_s now must denote the mass enclosed *at the start of the collapse*, while the new dimensionless variable $\mu(m_s, t)$ is the fraction of mass at a coordinate m_s which has not yet annihilated at the later time t . Then the density ρ is simply

$$\rho(m_s, t) = \frac{\mu(m_s, t)}{4\pi r^2(m_s, t) \partial r(m_s, t) / \partial m_s} . \quad (53)$$

In the presence of annihilation, the collapse equation becomes

$$\frac{\partial^2 r(m_s, t)}{\partial t^2} = - \frac{G}{r^2(m_s, t)} \int_0^{m_s} dm'_s \mu(m'_s, t) . \quad (54)$$

B. Numerical calculation

To solve our system of coupled partial differential equations numerically, we divided the collapsing sphere into N shells, initially with equal masses and with the n th such shell having a time-dependent mass $\Delta m_n(t)$ and radius $r_n(t)$. If $n > 1$, then the spatially discretized equations of motion for this shell are

$$\begin{aligned} \frac{d^2 r_n(t)}{dt^2} &= - \frac{G}{r_n(t)^2} \sum_{i=1}^n \Delta m_i(t) \\ \frac{d \Delta m_n(t)}{dt} &= - \frac{3 \langle \sigma v \rangle_{ann}}{4\pi m} \Delta m_n(t) \left(\frac{dm_s}{dr^3} \right)_n \\ &\approx - \frac{3 \langle \sigma v \rangle_{ann}}{4\pi m} \left(\frac{\Delta m_n(t)^2}{r_n(t)^3 - r_{n-1}(t)^3} \right) , \end{aligned} \quad (55)$$

where the central “shell” ($n = 1$) is treated as a homogenous sphere as in the previous section, and thus will have the following equations of motion:

$$\begin{aligned} \frac{d^2 r_1(t)}{dt^2} &= - \frac{G}{r_1(t)^2} \Delta m_1(t) \\ \frac{d \Delta m_1(t)}{dt} &= - \frac{3 \langle \sigma v \rangle_{ann}}{4\pi m} \left(\frac{\Delta m_1(t)^2}{r_1(t)^3} \right) . \end{aligned} \quad (56)$$

Our requirement in the previous subsection that α be large also requires that this innermost shell is significantly larger than its annihilation radius at $t = 0$. For $\alpha = 3/4\pi \times 10^6$, $r_1(t = 0)$ is one hundred times greater than the annihilation radius for a homogenous sphere of mass $\Delta m_1(t = 0)$.

In order to nondimensionalize this system, we choose convenient units as before with the homogenous case; let the mass unit be the initial shell mass $\Delta m_0 = M_0/N$ and let the length unit be the previously-defined annihilation length for the central shell

$$R_{ann} = \left(\frac{3\langle\sigma v\rangle_{ann}}{4\pi m G} \right)^{2/3} \left(\frac{GM_0}{N} \right)^{1/3}. \quad (57)$$

We also choose a corresponding time unit

$$t_c = \sqrt{\frac{R_{ann}^3}{G\Delta m_0}} \quad (58)$$

such that the equations take on a simpler form in terms of the new dimensionless variables $\Delta\hat{m}_n = \Delta m_n/\Delta m_0$, $\hat{r}_n = r_n/R_{ann}$ and $\tau = t/t_c$:

$$\begin{aligned} \frac{d^2\hat{r}_n(\tau)}{d\tau^2} &= -\frac{1}{\hat{r}_n(\tau)^2} \sum_{i=1}^n \Delta\hat{m}_i(\tau) \\ \frac{d\Delta\hat{m}_n(\tau)}{d\tau} &\approx -\frac{\Delta\hat{m}_n(\tau)^2}{\hat{r}_n(\tau)^3 - \hat{r}_{n-1}(\tau)^3} \end{aligned} \quad (59)$$

for $n > 1$ and

$$\begin{aligned} \frac{d^2\hat{r}_1(\tau)}{d\tau^2} &= -\frac{\Delta\hat{m}_1(\tau)}{\hat{r}_1(\tau)^2} \\ \frac{d\Delta\hat{m}_1(\tau)}{d\tau} &\approx -\frac{\Delta\hat{m}_1(\tau)^2}{\hat{r}_1(\tau)^3} \end{aligned} \quad (60)$$

for $n = 1$. These equations were integrated in time using a fourth order Runge-Kutta routine similar to that in the above section, with the initial time $\tau = 0$, initial shell masses $\Delta\hat{m}_n = 1$ and initial radii given by equation (36) to be

$$\hat{r}_n(\tau = 0) = \left(\frac{4\pi\alpha n}{3} \right)^{1/3} \left(\gamma \left(\frac{n}{N} \right)^b + 1 \right)^{2/3}. \quad (61)$$

The initial infall velocities were chosen to be

$$\left(\frac{d\hat{r}_n}{d\tau} \right)_{\tau=0} = -\sqrt{\frac{2}{\hat{r}_n} \sum_{i=1}^n \Delta\hat{m}_i}, \quad (62)$$

which corresponds to zero energy infall.

As we would expect from the homogeneous case, each shell reached both zero size and zero mass at the same time in the absence of black hole formation, i.e. for clouds which annihilate completely. Some results for this simulation are shown in Figure 3 for $b = 2$, $\alpha = 3/4\pi \times 10^6$, $\beta_i = 100$ and $N = 1000$. We chose to use one thousand shells for the simulation because this N provided sufficient spatial resolution while still allowing the code to run within a reasonable amount of time.

It is important to note that, for this set of parameters, there appears to be a transition around a time $\tau \sim 500$. This is simply the annihilation time t_{ann} , defined in equation (46) to be the time at which the center reaches the annihilation density:

$$t_{ann} = t_0 - \frac{1}{\sqrt{6\pi G \rho_{ann}}} = t_0 \left(1 - \sqrt{\frac{3}{8\pi\alpha}} \right); \quad (63)$$

using $t_0 = \sqrt{\alpha/6\pi}(\langle\sigma v\rangle_{ann}/Gm)$, $\alpha = 3/4\pi \times 10^6$ and our nondimensional time variable, we find that $\tau_{ann} \equiv t_{ann}/t_c \approx 471$. This is the same value as we see in the numerical solution, and therefore we can conclude that neglecting annihilation effects until reaching the critical length scale is a good approximation for this choice of parameters.




FIG. 3: Data for a simulation run in which a black hole does not form, with $b = 2$, $N = 1000$, $\alpha = 3/4\pi \times 10^6$ and $\beta = 100$. The top plot shows the log of the dimensionless density $\hat{\rho}$ versus the log of the radius $\hat{r}$ for several times $\tau = 0, 200, 500, 1000$ and 2000. The bottom plot shows the dimensionless mass enclosed versus radius for the same τ values.

C. Black hole formation

Our main goal is to generalize our investigation of black hole formation in uniform density collapse to non-uniform collapse. To that end, at each time step, we check to see if any shell satisfies the condition

$$\frac{2Gm_s}{r} \geq 1 ; \quad (64)$$

in terms of our nondimensional variables and spatial discretization, this condition becomes

$$\left[\frac{1}{2N^2} \left(\frac{M_0}{M_{ann}} \right)^2 \right]^{1/3} \left(\frac{1}{\hat{r}_n} \sum_{n=1}^i \Delta \hat{m}_i \right) \geq 1 . \quad (65)$$

If this condition is true for a given shell, then we stop integrating the equations of motion for all shells inside it. In other words, we find the location of the event horizon and shut off the annihilation and collapse of the mass inside it. Then, if a black hole does form, there will be an initial seed hole that subsequently will accrete mass from the remaining (annihilating) matter outside. If a black hole does not form, then all of the shells will eventually annihilate away until no mass is left. Again, we shall see that the initial cloud mass M_0 must be above some critical minimum value, M_c , in order for the cloud to collapse to a black hole.

In general, the mass scale M_c depends on β_{ann} and b . To determine M_c for a given choice of b and β_{ann} , we increased our inputted value of the mass ratio M_0/M_{ann} until a black hole could form. (Of course, $\beta_{ann} = 1$ reproduced the results found for the homogenous case, independent of the value of b .) As we increased the contrast β_{ann} , we found that this minimum mass, or M_c , also increased via a rough power law. These results, for several b values, are plotted in Figure 4. Note that the exponent of the power law depends on b .

We can estimate the critical masses in Figure 4 by using the LL solution and simultaneously requiring that a black hole forms before the annihilation time t_{ann} . In the LL solution,

$$\phi \equiv \frac{2Gm_s}{r} = \left[\frac{4Gm_s}{3(Bm_s^b + t_0 - t)} \right]^{2/3} , \quad (66)$$




FIG. 4: Simulation data plot of $\log(M_c/M_{ann})$ versus $\log(\beta_{ann})$ for $b = 0.1, 1$ and 2 .

and this has a maximum value when

$$m_s = \left(\frac{t_0 - t}{B(b-1)} \right)^{1/b}; \quad (67)$$

note that this becomes singular as $b \rightarrow 1$, and therefore we will assume $b > 1$ for the time being. Inserting equation (67) into our equation for ϕ gives us the maximum value

$$\phi_{max} \equiv \left[\frac{4G(b-1)^{1-1/b}}{3bB^{1/b}(t_0-t)^{1-1/b}} \right]^{2/3}; \quad (68)$$

$\phi_{max} = 1$ at the time of initial black hole formation, t_{BH} , i.e. when

$$t = t_{BH} \equiv t_0 - \left(\frac{4G}{3} \right)^{\frac{b}{b-1}} \left(\frac{b-1}{B^{\frac{1}{b-1}} b^{\frac{b}{b-1}}} \right). \quad (69)$$

In order to produce a black hole before annihilation becomes important, we want $\phi_{max} = 1$ and $t < t_{ann}$; this happens if

$$B < \left(\frac{4G}{3} \right)^b \frac{(b-1)^{b-1}}{b^b (t_0 - t_{ann})^{b-1}}. \quad (70)$$

Using equation (48), we can rewrite this condition as

$$M_0 > \frac{3b(t_0 - t_{ann})}{4G(b-1)^{1-1/b}} \left[\sqrt{\left(\frac{b+1}{2b+1} \right)^2 + \frac{\beta_{ann}-1}{2b+1}} - \left(\frac{b+1}{2b+1} \right) \right]^{1/b} \equiv M_c. \quad (71)$$

For large β_{ann} , this yields the power law $M_c \propto \beta_{ann}^{1/2b}$.

If $b \leq 1$, the situation is somewhat different. Note that, for any b value,

$$\frac{\partial}{\partial m_s} \left(\frac{2Gm_s}{r} \right) \propto t_0 - t + (1-b) B m_s^b, \quad (72)$$

which is positive for $b \leq 1$ and $t_0 - t > 0$, indicating that ϕ is an increasing function of m_s under these circumstances. In that case, ϕ is largest at the edge, where $m_s = M_0$. Following the same line of reasoning as for the $b > 1$ case, requiring that $\phi_{max} = 1$ and $t < t_{ann}$, we find that

$$M_c = \frac{3(t_0 - t_{ann})}{4G} \left(\sqrt{\left(\frac{b+1}{2b+1}\right)^2 + \frac{\beta_{ann} - 1}{2b+1}} + \frac{b}{2b+1} \right) \quad (73)$$

for the $b \leq 1$ case. Using the data in Figure 4, we have found that this estimate is too large by a factor ~ 2 for the $b = 0.1$ case, even though the data do follow the predicted power law $M_c \propto \beta_{ann}^{1/2}$ for larger values of β_{ann} ; in general, the estimate of equation (73) is too large by factors of order unity for b values less than one. Hence, for $b \leq 1$, we can conclude that it is possible for black holes to form at a later time $t = t_{ann} + \Delta t$, such that M_c is reduced by a factor of $(t_0 - t_{ann} - \Delta t)/(t_0 - t_{ann})$.

The simulation also tracked the masses of both the initial seed black holes and the end-state black holes that they eventually evolved to. Theoretically, for $b \leq 1$, we expect that the initial black hole will have a mass of order M_0 , and this prediction was verified by the simulation. For $b > 1$, we should find the initial black hole mass from equation (67):

$$M_{BH} = \left(\frac{t_0 - t_{BH}}{B(b-1)} \right)^{1/b} ; \quad (74)$$

plugging in what we know for t_{BH} gives us

$$M_{BH} = \left(\frac{M_0^b}{bM_{ann}} \left[\sqrt{\left(\frac{b+1}{2b+1}\right)^2 + \frac{\beta_{ann} - 1}{2b+1}} - \frac{b+1}{2b+1} \right]^{-1} \right)^{\frac{1}{b-1}} \propto M_{ann} \left(\frac{M_0}{M_{ann}} \right)^{\frac{b}{b-1}} \quad (75)$$

which agrees with the simulation data.

Finding the final black hole mass is equivalent to finding the total mass change due to annihilation, which we will call ΔM . Suppose that a black hole forms with a mass that is M_{BH} initially. The mass density at this time is $\rho_{BH} \sim 3/32\pi G^3 M_{BH}^2$, so annihilation proceeds at a rate $\approx \rho_{BH} \langle \sigma v \rangle_{ann} M_{BH} / m \sim 3 \langle \sigma v \rangle_{ann} / 32\pi G^3 m M_{BH} \sim M_{ann} / G M_{BH}$. Up to numerical factors, we therefore expect a peak annihilation luminosity $-\dot{M} \sim M_{ann} / G M_{BH}$ and a total mass loss to annihilation $\Delta M \sim -M_{ann}$. Thus, whenever a black hole forms, a similar amount of energy is released in the form of annihilation products.

To study this more completely, we consider collapsing clouds well above the critical mass for black hole formation. The easiest case is the collapse of a homogenous cloud ($b = 0$). If we assume that the initial mass is significantly above M_c , then we can assume that annihilation will not affect the trajectories of the shells as they fall inward. For the uniform case

$$\rho(t) = \frac{1}{6\pi G (t_0 - t)^2} \quad (76)$$

and a black hole forms when $t_0 - t_{BH} = 4GM_0/3$. The total mass annihilated is

$$\begin{aligned} \Delta M &= -\frac{\langle \sigma v \rangle_{ann} M_0}{6\pi G m} \int_{-\infty}^{t_{BH}} \frac{dt}{(t_0 - t)^2} = -\frac{\langle \sigma v \rangle_{ann} M_0}{6\pi G m} \left(\frac{1}{t_0 - t_{BH}} \right) \\ &= -\frac{\langle \sigma v \rangle_{ann}}{8\pi G^2 m} = -\frac{2}{3} M_{ann} . \end{aligned} \quad (77)$$

Therefore, the total mass that is lost to annihilation in the homogenous case is of order M_{ann} , for any initial mass that is greater than the minimum mass M_c .

All cases with $b \leq 1$ are similar to the $b = 0$ case in that the event horizon first forms at the outer edge. In this case, annihilation ends when

$$t_0 - t_{BH} = \frac{4M_0}{3} - B M_0^b ; \quad (78)$$

using the LL solution for $\rho(m_s, t)$ we find

$$\Delta M = -\frac{\langle \sigma v \rangle_{ann}}{12\pi b G m B M_0^{b-1}} \int_0^1 \frac{dx}{x^b} \ln \left[\frac{(2b+1) B M_0^b x^b - B M_0^b + 4M_0/3}{B M_0^b x^b - B M_0^b + 4M_0/3} \right] . \quad (79)$$

In the limit that b goes to zero, equation (79) yields the same answer as before: $\Delta M \rightarrow -2M_{ann}/3$.

For $b > 1$, the situation is more complicated because annihilation continues after the initial black hole formation. For this case, it is useful to think in terms of the luminosity as a function of time, which is

$$\begin{aligned} L(t) \equiv -\frac{d\Delta M}{dt} &= \frac{\langle \sigma v \rangle_{ann}}{m} \int_{M_{min}(t)}^{M_0} \rho(m_s, t) dm_s \\ &= \frac{\langle \sigma v \rangle_{ann}}{6\pi Gm} \int_{M_{min}(t)}^{M_0} \frac{dm_s}{(Bm_s^b + t_0 - t) [(2b+1) Bm_s^b + t_0 - t]} , \end{aligned} \quad (80)$$

where $M_{min}(t)$ is the mass within the event horizon, which doesn't contribute to the luminosity. Let t_{BH} be the time when a black hole first forms; then for $t < t_{BH}$, we have $M_{min}(t) = 0$, and for $t > t_{BH}$ we have

$$\frac{4GM_{min}(t)}{3} = B [M_{min}(t)]^b + t_0 - t . \quad (81)$$

For very early times, when $t < t_{BH}$ and $t_0 - t \gg Bm_0^b$, there is an initial phase where we can approximate

$$\begin{aligned} \left(\frac{d\Delta M}{dt} \right)_{early} &= -\frac{\langle \sigma v \rangle_{ann}}{6\pi Gm} \int_0^{M_0} \frac{dm_s}{(Bm_s^b + t_0 - t) [(2b+1) Bm_s^b + t_0 - t]} \\ &\approx -\frac{\langle \sigma v \rangle_{ann}}{6\pi m} \left[\frac{M_0}{(t_0 - t)^2} \right] . \end{aligned} \quad (82)$$

For smaller $t_0 - t$, but still before black hole formation, we have a another phase where

$$\left(\frac{d\Delta M}{dt} \right)_{t < t_{BH}} = -\frac{\langle \sigma v \rangle_{ann}}{6\pi Gm (t_0 - t)^2} \int_0^{M_0} \frac{dm_s}{[Bm_s^b / (t_0 - t) + 1] [(2b+1) Bm_s^b / (t_0 - t) + 1]} ; \quad (83)$$

if we define $x^b \equiv Bm_s^b / (t_0 - t)$ and assume that $(M_0/M_{BH})^b \gg 1$, then this becomes

$$\left(\frac{d\Delta M}{dt} \right)_{t < t_{BH}} \approx -\frac{b^2 M_{ann}}{2(b-1)^{2-1/b} GM_{BH}} \left(\frac{t_0 - t_{BH}}{t_0 - t} \right)^{2-1/b} \int_0^\infty \frac{dx}{(x^b + 1) [(2b+1) x^b + 1]} . \quad (84)$$

The $d\Delta M/dt$ of this second phase increases with time until it ends right before $t = t_{BH}$, and so we get a peak luminosity of

$$\dot{M}_{peak} \approx \frac{b^2 M_{ann}}{2(b-1)^{2-1/b} GM_{BH}} \int_0^\infty \frac{dx}{(x^b + 1) [(2b+1) x^b + 1]} . \quad (85)$$

Right at $t = t_{BH}$, we know that $M_{min} = M_{BH}$, and so then we have

$$\left(\frac{d\Delta M}{dt} \right)_{t=t_{BH}} \approx -\frac{b^2 M_{ann}}{2(b-1)^{2-1/b} GM_{BH}} \int_{(b-1)^{-1/b}}^\infty \frac{dx}{(x^b + 1) [(2b+1) x^b + 1]} , \quad (86)$$

and thus, in our simple model, $d\Delta M/dt$ drops by a factor of order one right after a black hole forms.

The total mass that is annihilated before t_{BH} , which we call ΔM_1 , is simply

$$\Delta M_1 = \int_{-\infty}^{t_{BH}} dt \frac{d\Delta M(t)}{dt} ; \quad (87)$$

the mass annihilated from the phase where $d\Delta M/dt \approx (d\Delta M/dt)_{early}$ is approximately

$$\Delta M_{early} \approx -\frac{2bM_{ann}}{3} \left(\frac{M_{BH}}{M_0} \right)^{b-1} \quad (88)$$

which is very small for $M_0/M_{BH} \gg 1$, and therefore we will neglect this term; then equation (87) becomes

$$\Delta M_1 \approx \int_{-\infty}^{t_{BH}} dt \left(\frac{d\Delta M(t)}{dt} \right)_{t < t_{BH}} \approx -\frac{2b^2 M_{ann}}{3(b-1)^{2-1/b}} \int_0^\infty \frac{dx}{(x^b + 1) [(2b+1) x^b + 1]} , \quad (89)$$

where we took the upper mass limit to infinity, which is justified for $M_0 \gg M_{BH}$ and $b > 1$.

After t_{BH} , we need to use the more general equations (80) and (81), but for very late times, when $t - t_0 \gg GM_{min}(t)$, we can approximate

$$M_{min}(t) \approx \left(\frac{t - t_0}{B} \right)^{1/b} (1 + \epsilon) , \quad (90)$$

where

$$\epsilon = \frac{4M_{min}(t)}{3b(t - t_0)} ; \quad (91)$$

then, in this very late phase, we can define the integration variable $x \equiv m_s/M_{min}(t)$ to get

$$\begin{aligned} \left(\frac{d\Delta M}{dt} \right)_{late} &\approx - \frac{\langle \sigma v \rangle_{ann} M_{min}(t)}{6\pi Gm(t - t_0)^2} \int_1^\infty \frac{dx}{[(1 + \epsilon)x^b - 1][(2b + 1)x^b - 1]} \\ &\approx - \frac{b^2 M_{ann}}{2(b - 1)^{1-1/b} GM_{BH}} \left(\frac{t_0 - t_{BH}}{t - t_0} \right)^{2-1/b} \ln \left(\frac{t - t_0}{t_0 - t_{BH}} \right) . \end{aligned} \quad (92)$$

The total mass annihilated after t_{BH} , which we call ΔM_2 , is given by

$$\Delta M_2 = - \frac{\langle \sigma v \rangle_{ann}}{6\pi Gm} \int_{M_{BH}}^{M_0} dm_s \int_{t_{BH}}^{t_f(m_s)} \frac{dt}{(Bm_s^b + t_0 - t)[(2b + 1)Bm_s^b + t_0 - t]} , \quad (93)$$

where $t_f(m_s)$ is the time at which a mass shell with a coordinate m_s reaches the event horizon:

$$\frac{4Gm_s}{3} = Bm_s^b + t_0 - t_f(m_s) ; \quad (94)$$

if we use the integration variable $x \equiv m_s/M_{BH}$, we find that

$$\frac{\Delta M_2}{M_{ann}} \approx - \frac{1}{3} \int_1^\infty \frac{dx}{x^b} \ln \left\{ (2x^{b-1} + 1) \left[\frac{(2b + 1)x^b + b - 1}{x^b + b - 1} \right] \right\} \quad (95)$$

for $M_0 \gg M_{BH}$ and $b > 1$.

We can integrate the above equations numerically to get ΔM_1 , ΔM_2 and $\dot{M}_{peak}$, and we show the results for several values of b in Table I. We also provide a sample lightcurve in Figure 5, which shows the scaled luminosity

TABLE I: Data for several values of b

b	$\Delta M_1/M_{ann}$	$\Delta M_2/M_{ann}$	$G\dot{M}_{peak}M_{BH}/M_{ann}$
2	-1.29	-1.08	-0.97
3	-1.01	-0.57	-0.76
4	-0.91	-0.39	-0.68


LM_{BH}/M_{ann} as a function of the scaled time $t(s)(10^7 M_\odot/M_{BH}) \approx (50 \text{ s})(t/GM_{BH})$. Our calculated light curve ignores the effects of opacity and general relativity. The former should be relatively unimportant when a black hole forms, as was discussed for the uniform case in Section IIB. (When $b < 1$, the uniform case is a good approximation, and when $b > 1$, most of the optical depth for annihilation products is near the hole.) General relativity would tend to lower the luminosity peak and smooth it out. Within factors of ~ 2 or 3 , we still expect a lightcurve resembling Figure 5, and a total annihilated mass $\sim M_{ann}$.

IV. DISCUSSION

A. P-Wave Annihilation

P-wave annihilation has $\langle \sigma v \rangle_{ann} = \sigma_0 \langle v^2 \rangle$. In the early Universe, $\langle v^2 \rangle = T/m$, and following [16] we find

$$\sigma_0 \approx 1.7 \times 10^{-7} \text{ GeV}^{-2} \left(\frac{x_f^2}{10^3 \Omega_{CDM} h^2 g_{\star S} / \sqrt{g_{\star}}} \right) , \quad (96)$$

FIG. 5: Plot of $L(t)$ versus t for $b = 2$.

where $x_f \approx C - \frac{3}{2} \ln(\ln C)$ with $C \approx 0.076(g/\sqrt{g_*})m\sigma_0/\sqrt{G}$. Assuming that thermal particles scatter from the CDM particles with a cross-section $f_t\sigma_0(T/m)^2$, we find that the CDM remains in kinetic equilibrium until

$$T_K \approx 1.6 \times 10^{-3} \text{ GeV} \frac{g_{*S}^{1/4} [m(\text{GeV})]^{3/4}}{g_t^{1/4} f_t^{1/4}} \left(\frac{10^3 \Omega_{\text{CDM}} h^2}{x_f^2} \right)^{1/4}; \quad (97)$$

consequently

$$K_{\text{CDM}} \approx 10^9 \text{ GeV}^{-8/3} \frac{g_t^{1/4} f_t^{1/4}}{[m(\text{GeV})]^{7/4} g_{*S}^{1/4} g_{*SK}^{2/3}} \left(\frac{x_f^2}{10^3 \Omega_{\text{CDM}} h^2} \right)^{1/4} \left(\frac{0.15}{\Omega_{\text{CDM}} h^2} \right)^{2/3}. \quad (98)$$

As long as the collapse proceeds adiabatically, $\langle v^2 \rangle = K_{\text{CDM}} \rho^{2/3}$, and for pure p-wave annihilation we find

$$M_{\text{ann}} \approx \frac{2.4 \times 10^8 M_\odot}{[m(\text{GeV})]^{33/28}} \left(\frac{g_t^{3/28} f_t^{3/28} g_{*S}^{13/28}}{g_{*S}^{15/28} g_{*SK}^{2/7}} \right) \left(\frac{x_f^2}{10^3 \Omega_{\text{CDM}} h^2} \right)^{15/28} \left(\frac{0.15}{\Omega_{\text{CDM}} h^2} \right)^{2/7}. \quad (99)$$

For example, if $m = 50 \text{ GeV}$ and $g_t \approx g_* \approx g_{*S} \approx g_{*SK} \approx 50$, we find $M_{\text{ann}} \approx 10^6 f_t^{3/28} M_\odot$. Not suprisingly, the value of M_{ann} is considerably smaller than for s-wave annihilation, under the assumption of adiabatic collapse: p-wave annihilation is less efficient, and therefore the cloud must reach higher density to annihilate, which lowers the critical mass for black hole formation.

In actuality, we do not expect the cloud to collapse adiabatically, and the critical mass is closer to the value for s-wave annihilation at $\langle \sigma v \rangle_{\text{ann}} = \sigma_0$. The reason is that ρ_{ann} is large enough that clouds become opaque to their annihilation products *before* they either annihilate or collapse catastrophically. Defining $\rho_{1,\text{ad}}$ to be the density at which the optical depth is one in adiabatic collapse, we find

$$\rho_{1,\text{ad}} = \rho_{\text{ann}} f_s^{-3/2} (K \rho_{\text{ann}}^{2/3})^{3/2} \left(\frac{M_{\text{ann}}}{M} \right)^{1/2} = \rho_{\text{BH}} f_s^{-3/2} (K \rho_{\text{ann}}^{2/3})^{3/2} \left(\frac{M}{M_{\text{ann}}} \right)^{3/2}; \quad (100)$$

since

$$K_{\text{CDM}} \rho_{\text{ann}}^{2/3} \approx \frac{1.2 \times 10^{-3}}{[m(\text{GeV})]^{5/28}} \left(\frac{g_t^{3/28} f_t^{3/28} g_{*S}^{13/28}}{g_{*S}^{15/28} g_{*SK}^{2/7}} \right) \left(\frac{10^3 \Omega_{\text{CDM}} h^2}{x_f^2} \right)^{13/28} \left(\frac{0.15}{\Omega_{\text{CDM}} h^2} \right)^{2/7}, \quad (101)$$

we see that the clouds become opaque before reaching their final destinies.

Once the clouds become opaque, they start to heat up. To understand what happens qualitatively, we introduce new density and mass scales

$$\begin{aligned}\rho_1 &= \frac{8\pi G m^2}{3f_s^2 \sigma_0^2} = \rho_{\text{ann}} \left(\frac{K \rho_{\text{ann}}^{2/3}}{f_s} \right)^2 \\ M_1 &= \sqrt{\frac{3}{32\pi G^3 \rho_1}} = \frac{3f_s \sigma_0}{16\pi G^2 m} = M_{\text{ann}} \left(\frac{f_s}{K \rho_{\text{ann}}^{2/3}} \right),\end{aligned}\quad (102)$$

in terms of which we find

$$\begin{aligned}\tau &= \left(\frac{\rho}{\rho_1} \right)^{2/3} \left(\frac{M}{M_1} \right)^{1/3} \\ v_{\text{ff}}^2 &= \frac{2GM}{R} = \left(\frac{\rho}{\rho_1} \right)^{1/3} \left(\frac{M}{M_1} \right)^{2/3};\end{aligned}\quad (103)$$

we can eliminate the density from these equations to find

$$v_{\text{ff}}^2 = \tau^{1/2} \left(\frac{M}{M_1} \right)^{1/2}. \quad (104)$$

Thus, we see that clouds with $M \geq M_1$ collapse to black holes before becoming opaque, and vice-versa. We also see that during adiabatic collapse,

$$\begin{aligned}\frac{\Delta M}{M} &\equiv \frac{\Gamma_{\text{ann}}}{H} = \left(\frac{K \rho_{\text{ann}}^{2/3}}{f_s} \right)^{7/3} \tau^{7/4} \left(\frac{M_1}{M} \right)^{7/12} \\ K \rho^{2/3} &= f_s \left(\frac{K \rho_{\text{ann}}^{2/3}}{f_s} \right)^{7/3} \tau \left(\frac{M_1}{M} \right)^{1/3} = \frac{f_s}{\tau^{3/4}} \left(\frac{M}{M_1} \right)^{1/4} \frac{\Delta M}{M}.\end{aligned}\quad (105)$$

Note that $\Delta M/M$ is the fractional mass annihilated in one free-fall time up to a factor of order unity.

Consider the fate of a cloud with $M/M_1 \lesssim 1$, but not $\ll 1$ (we shall estimate how much smaller shortly). Once the cloud reaches $\tau > 1$, the relativistic products of the annihilation do not escape freely, and deposit a substantial fraction of their total energy into kinetic energy of the massive particles. Because the optical depth for the relativistic particles is proportional to $(E/m)^2$, after a relativistic particle scatters once, thereby losing about half of its original energy on average, its optical depth to escape drops by a substantial factor; hence, as long as τ is not too large, relativistic particles are not trapped. The massive particles that scatter the relativistic particles are heated to semi-relativistic kinetic energies i.e. $\sim m/2$. For these, however, the scattering cross section remains $\approx \sigma_0$; moreover, on each scatter off of one of the colder massive particles that dominate the cloud, these hot massive particles lose of order half of their kinetic energy. As a result, they continue to slow down, and it is likely that they become trapped inside the cloud. The second of equations (105) shows that $\Delta M/M \gtrsim K \rho^{2/3}$ for clouds masses $\lesssim M_1$ at $\tau \gtrsim 1$, so the cloud heats up to $\langle v^2 \rangle = \eta \Delta M/M$, with $\eta \lesssim 1/2$. Thus, we expect the internal energy of the cloud to rise rapidly to $\eta \Delta M/M$ as it continues to collapse to larger optical depth.

Since the internal energy of the cloud rises as it annihilates, the annihilation equation is altered to

$$\frac{dM}{dt} = -\frac{d\Delta M}{dt} \approx -\frac{\eta \rho \sigma_0 \Delta M}{m}; \quad (106)$$

for $\Delta M \lesssim M$ we find

$$\ln \Delta M = \ln \Delta M_i + \frac{2\eta}{3} \left[\left(\frac{\rho}{\rho_1} \right)^{1/2} - \left(\frac{\rho_i}{\rho_1} \right)^{1/2} \right]. \quad (107)$$

Taking ρ_i and M_i to correspond to $\tau = 1$, we find that $\Delta M \rightarrow M$ when the density must increase to

$$\left(\frac{\rho}{\rho_1} \right)^{1/2} \approx \frac{7}{2\eta} \ln \left(\frac{f_s}{K \rho_{\text{ann}}^{2/3}} \right) - \frac{7}{8\eta} \ln \left(\frac{M_1}{M} \right) + \left(\frac{M_1}{M} \right)^{1/4}, \quad (108)$$

for the cloud to annihilate away. (The last stages of the cloud's collapse could deviate somewhat from free-fall, as the pressure builds up to values comparable to ρv_{ff}^2 , but this phase most likely is only short-lived.) Thus, we estimate that the critical mass for a black hole to form is of order

$$M_{c,p} = \frac{2\eta M_1}{7 \ln(f_s/K\rho_{\text{ann}}^{2/3})} = \frac{2\eta f_s M_{\text{ann}}}{7 K \rho_{\text{ann}}^{2/3} \ln(f_s/K\rho_{\text{ann}}^{2/3})}; \quad (109)$$

for example, if $m = 50$ GeV and $g_t \approx g_\star \approx g_{\star S} \approx g_{\star SK} \approx 50$, we find that $M_{c,p} \approx 10(2\eta)M_{\text{ann}}$ or of order $10^7(2\eta)M_\odot$. We note that since M_1 is of order M_{ann} for s-wave annihilation with $\langle\sigma v\rangle_{\text{ann}} = \sigma_0$, the critical mass for black hole formation for p-wave annihilation is only smaller by a factor of $\sim 7 \ln(f_s/K\rho_{\text{ann}}^{2/3})/2\eta$, not by the factor of $\sim K\rho_{\text{ann}}^{2/3}$ deduced under the false assumption of adiabatic collapse.

It is also possible that the CDM particles can annihilate via both s-wave and p-wave channels with p-wave dominant in the early Universe but s-wave dominant during the early stages of the cloud collapse. If the s-wave annihilation has $\langle\sigma v\rangle_{\text{ann}} = \epsilon_s \sigma_0$, then it dominates until $\Delta M/M \approx \epsilon_s$, whereupon the p-wave channel takes over. In this situation, we expect to get a critical mass for black hole formation that is smaller than for s-wave annihilation with $\langle\sigma v\rangle_{\text{ann}} = \sigma_0$ by a logarithmic factor of order $(3/2\eta) \ln \epsilon_s^{-1}$.

B. Conclusions

Our calculations show that the collapse of a cloud of annihilating matter to a black hole is only possible for masses above a critical mass, M_c , that is of order $10^7 - 10^8 M_\odot$ for plausible cold dark matter candidates. For a radial density profile, the critical mass depends on the parameters of the profile; for example, we found that $M_c/M_{\text{ann}} \sim \beta_{\text{ann}}^{1/2b}$ for an LL profile with the exponent b [23], where β_{ann} is the density contrast between the center and the edge when annihilation becomes important. The conditions for collapse studied here are highly idealized since we assumed that the clouds are exactly spherical and cold, at least initially. Internal shearing motions induced by tidal perturbations in the early Universe would violate these assumptions. However, it is conceivable that the collapse of larger, sheared structures could very rarely lead to the formation of bound fragments similar to what we have assumed here [19, 20]. In any event, it is likely that we have assumed conditions most favorable for forming a black hole from CDM collapse, so our lower bounds on the masses of black holes that can form in this way are probably, if anything, underestimates of reality. It is remarkable, although perhaps completely fortuitous, that the masses estimated here – which are determined by fundamental constants of Nature – are even close to the sorts of black hole masses observed astronomically [17].

When a cloud collapses to a black hole, annihilation releases energy $\sim M_{\text{ann}}$ in a time $\sim GM_{BH}$, where $M_{BH} > M_{\text{ann}}$ is the mass of the black hole when collapse first occurs. The peak luminosity resulting from the annihilation is therefore $\sim M_{\text{ann}}/GM_{BH}$, and much of the annihilation occurs in a prodigious, relatively short-lived burst. Thus the formation of a black hole in the collapse of annihilating CDM would be a spectacular event: for s-wave annihilation, approximately $M_{\text{ann}} \sim 2 \times 10^{64} [m(\text{GeV})]^{-1} (x_f \sqrt{g_\star}/15g_{\star S})$ ergs would be released in a time of order $10^5 [m(\text{GeV})]^{-1} (x_f \sqrt{g_\star}/15g_{\star S}) (M_{BH}/M_{\text{ann}})$ seconds, with a peak luminosity of $4 \times 10^{59} (M_{\text{ann}}/M_{BH})$ ergs per second; for $m = 50$ GeV, the energy released is of order 7×10^{61} ergs over about $380 (M_{BH}/M_{\text{ann}})$ seconds. For comparison, note that $\frac{1}{2}(10^{11} M_\odot)(300 \text{ km s}^{-1})^2 \approx G(10^{11} M_\odot)^2/10 \text{ kpc} \approx 10^{59}$ ergs; i.e. the energy released in forming a single annihilating CDM black hole is comparable to, if not larger than, the binding energy of an entire galaxy.

What an observer could detect from such events depends on what annihilation products form, but even if the efficiency of generating electromagnetic radiation is $\lesssim 10^{-8}$ of the total energy released, they would be as bright as extragalactic gamma ray bursts. Any direct production of standard model particles in the annihilation should produce a pair fireball with little or no net baryon content, even for extremely low branching ratios for $mm \rightarrow$ (standard model pairs). For example, if there is no direct production of $e^\pm$ pairs or quarks, but a fraction f_ν of the annihilation energy goes into neutrinos and antineutrinos of energy m , then we estimate an efficiency of producing $e^\pm$ pairs of about $(0.01 - 0.1) f_\nu G_F^2 m^2 / \langle\sigma v\rangle_{\text{ann}} \approx (0.001 - 0.01) f_\nu [m(\text{GeV})]^2 (x_f \sqrt{g_\star}/15g_{\star S})^{-1}$ for s-wave annihilation, and perhaps a bit smaller for p-wave annihilation; this would lead to a total energy of order $(0.02 - 0.002) \times 10^{64} f_\nu^2 m(\text{GeV})$ ergs [26]. Thus, a tiny branching ratio for $mm \rightarrow \nu\bar{\nu}$ would result in the formation of an extremely energetic pair fireball, even without any direct pair production via $mm \rightarrow e^+e^-$ or $mm \rightarrow q\bar{q}$: if $f_\nu \sim 10^{-5}$, we estimate a total pair fireball energy of order $10^{51} - 10^{52} m(\text{GeV})$ ergs. Acceptable branching ratios for direct production of e^+e^- or $q\bar{q}$ pairs are far smaller, $\sim 10^{-10}$. The latest upper bounds on the WIMP-nucleon scalar cross section from CDMS [25] are in the range $10^{-42.4} - 10^{-41} \text{ cm}^2 \approx 10^{-15} - 10^{-13.6} \text{ GeV}^{-2}$, compared to $\langle\sigma v\rangle_{\text{ann}} \approx 8.6 \times 10^{-9} \text{ GeV}^{-2} (x_f \sqrt{g_\star}/15g_{\star S})$ for s-wave annihilation and $\sigma_0 \approx 1.7 \times 10^{-7} \text{ GeV}^{-2} (x_f^2 \sqrt{g_\star}/150g_{\star S})$ for p-wave annihilation, so very low branching ratios for $mm \rightarrow$ (standard model pairs) may be reasonable, if not expected.

Astronomical observations could hardly have missed the production of pair fireballs with energy $\sim M_{ann}$ on timescales $\sim GM_{BH}$. Moreover, a blast wave composed of $e^\pm$ pairs, photons and baryons with a total energy that is even a modest fraction of M_{ann} (say 0.001 – 0.01) could deposit enough energy to substantially alter if not completely eject the interstellar medium of a galaxy. One possibility is that the branching ratios are $\lesssim 10^{-10}$ for $mm \rightarrow e^+e^-$ or $q\bar{q}$ and $\lesssim 10^{-5}$ for $mm \rightarrow \nu\bar{\nu}$, as discussed above. Alternatively, either black holes do not form from annihilating dark matter, or else the dark matter is in some other form that does not annihilate, such as axions.

Acknowledgments

R. A. Vanderveld has been supported by an NSF Graduate Research Fellowship and a Cornell NASA Space Grant Fellowship.

-
- [1] D. N. Spergel et al., *Astrophys. J. Suppl.* **148**, 175 (2003).
 - [2] J. R. Primack, *Nuc. Phys. B (Proc. Suppl.)* **124**, 3 (2003).
 - [3] B. W. Lee and S. Weinberg, *Phys. Rev. Lett.* **39**, 165 (1977).
 - [4] J. E. Gunn, B. W. Lee, I. Lerche, D. N. Schramm and G. Steigman, *Ap. J.* **223**, 1015 (1978).
 - [5] L. Bergström, *Rep. Prog. Phys.* **63**, 793 (2000).
 - [6] C. L. Bennett et al., *Astrophys. J. Suppl.* **148**, 1 (2003).
 - [7] R. Bean, A. Melchiorri and J. Silk, *Phys. Rev. D* **68**, 083501 (2003).
 - [8] M. Tegmark et al., *Phys. Rev. D* **69**, 103401 (2004).
 - [9] J. Dubinski and R. G. Carlberg, *Ap. J.* **378**, 496 (1991).
 - [10] J. F. Navarro, C. S. Frenk and S. D. M. White, *Ap. J.* **490**, 493 (1997).
 - [11] B. Moore, T. Quinn, F. Governato, J. Stadel and G. Lake, *Mon. Not. R. Astron. Soc.* **310**, 1147 (1999).
 - [12] M. Kaplinghat, L. Knox, M. S. Turner, *Phys. Rev. Lett.* **85**, 3335 (2000).
 - [13] S. Ghigna, B. Moore, F. Governato, G. Lake, T. Quinn and J. Stadel, *Ap. J.* **544**, 616 (2000).
 - [14] D. Merritt, M. Milosavljević, L. Verde and R. Jimenez, *Phys. Rev. Lett.* **88**, 191301 (2002).
 - [15] J. E. Taylor and J. Silk, *Mon. Not. R. Astron. Soc.* **339**, 505 (2003).
 - [16] E. W. Kolb and M. S. Turner, *The Early Universe*, (Addison-Wesley Publishing Company, 1990), equations 5.44 and 5.47, but using $T_{\gamma,0} = 2.73^\circ\text{K}$ instead of 2.75°K .
 - [17] J. Kormendy and D. Richstone, *Annu. Rev. Astron. Astrophys.* **33**, 581 (1995).
 - [18] P. J. E. Peebles, *Principles of Physical Cosmology*, (Princeton University Press, 1993).
 - [19] A. Loeb and F. A. Rasio, *Ap. J.* **432**, 52 (1994).
 - [20] A. Valinia, P. R. Shapiro, H. Martel and E. T. Vishniac, *Ap. J.* **479**, 46 (1997).
 - [21] W. H. Press, S. A. Teukolsky, W. T. Vetterling and B. P. Flannery, *Numerical Recipes in C*, 2nd ed. (Cambridge University Press, 1998), Ch. 16.
 - [22] M. V. Penston, *Mon. Not. R. Astron. Soc.* **144**, 425 (1969).
 - [23] D. Lynden-Bell and J. P. S. Lemos, *Mon. Not. R. Astron. Soc.* **233**, 197 (1988).
 - [24] S. N. Spergel and P. J. Steinhardt, *Phys. Rev. Lett.* **84**, 3760 (2000).
 - [25] D. S. Akerib et al., *Phys. Rev. Lett.* **93**, 211301 (2004).
 - [26] J. Goodman, A. Dar and S. Nussinov, *Ap. J.* **314**, L7 (1987).